

Entire Function

Next, we will discuss:

- The zeros of entire function
- How zeros determine an entire function.

(1) Jensen's Formula:

Thm. $D_{0,R} \subseteq \mathbb{C}$ open. $f \in \mathcal{O}(D_{0,R})$. $f(0) \neq 0$.

$f(z) \neq 0$. $\forall z \in D_{0,R}$. If $\{z_k\}_1^n$ seq
of zeros inside $D_{0,R}$. Then $\log |f(z)|$
 $= \sum_1^n \log \frac{|z_k|}{R} + \frac{1}{2\pi} \int_0^{2\pi} \log |f(Re^{i\theta})| d\theta$.

Pf. 1) Note that $g(z) = f(z) / \prod_{k=1}^n (z - z_k)$

Refining g at $\{z_k\}_1^n$, by series.

Then $g(z) \in \mathcal{O}(D_{0,R})$, $g(z) \neq 0$ in $D_{0,R}$

2) For $g(z)$: $\exists h(z) \in \mathcal{O}(D_{0,R})$, s.t.

$$g(z) = e^{h(z)} \quad \therefore |\log(Re^{i\theta})| = C$$

By Mean value of harmonic $h(z)$.

$$\therefore \frac{1}{2\pi} \int_0^{2\pi} \log |g(Re^{i\theta})| d\theta = \log |g(z)|$$

3) For $z - z_k$:

$$\text{prove: } \log |z_k| = \log \frac{|z_k|}{R} + \frac{1}{2\pi} \int_0^{2\pi} \log |Re^{i\theta} - z_k| d\theta$$

$$\text{Note } |Re^{i\theta} - z_k| = |R - z_k e^{-i\theta}| \neq 0.$$

Similar method of 2). We have

$$\text{mean value of } R - z_k e^{-i\theta}$$

$$4) \text{ Note that } \frac{i}{2\pi} \int_0^{2\pi} \log |f(re^{i\theta})| d\theta \\ = \frac{i}{2\pi} \int_0^{2\pi} \log |g(re^{i\theta})| d\theta + \frac{i}{2\pi} \sum_k^n \int_0^{2\pi} \log |re^{i\theta} - z_k| d\theta$$

Remark: From Jensen Formula, we can connect the growth of holomorphic $f(z)$ with its zeros number.

Def: $n(r)$ is the number of zeros of $f(z)$ which are inside $D(0, r)$.

$$\text{Cir.} \quad \int_0^r n(r) \frac{dr}{r} = \frac{i}{2\pi} \int_0^{2\pi} \log |f(re^{i\theta})| d\theta - \log |f(0)|.$$

$$\underline{\text{Pf:}} \quad \underline{\text{Lemma}} \quad \int_0^r n(r) \frac{dr}{r} = \sum_k^n \log \left| \frac{r}{z_k} \right|$$

$$\text{Note that } n(r) = \sum_k^n \chi_{\{|z_k| < r\}}$$

(2) Finite Orders:

$f \in \mathcal{O}(C)$. If there exists $\ell, A, B > 0$, s.t.

$|f(z)| \leq A e^{B|z|^\ell}$. Then we say the order of $f \leq \ell$. Def $\ell_f = \inf \ell$

Thm. For $f \in \mathcal{O}(C)$, $\ell_f \stackrel{a}{=} \ell$.

i) $n(r) \leq Cr^\ell$, for some $C > 0$, r is large enough

ii) $\{z_k\}_1^n$ seq of zeros of $f(z)$, $z_k \neq 0$, $\forall k \in \mathbb{Z}^+$.

Then $\forall s > \ell$, we have: $\sum \frac{1}{|z_k|^s} < \infty$

Remark: The number of zeros is restricted by the order of entire function.

Pf: i) For applying Jensen Formula:

Consider $F(z) = f(z)/z^k$, k is multiple of zeros $\neq 0$

$$\therefore n_F(r) = n_f(r) - k, \quad \ell_F = \ell_f.$$

$$\text{Note that } \int_0^R n(r) \frac{dr}{r} = \frac{1}{2\pi} \int_{-\pi}^{\pi} |F(re^{i\theta})| - b_F(\theta)| d\theta$$

We need $n(r)$ jump out of integration in LHS.

$$\text{By monotone of } n(r), \quad \int_0^R n(r) \frac{dr}{r} \geq \int_{\frac{R}{2}}^R n\left(\frac{R}{2}\right) \frac{dr}{r}.$$

ii) From i):

$$\begin{aligned} \sum_{|z_k| \geq 1} \frac{1}{|z_k|^s} &= \sum_k \sum_{j: s/2k \in 2^{j+1}} |z_k|^{-s} \leq \sum n(2^{j+1}) 2^{-sj} \\ &\leq \sum 2^{c(j+1)} \cdot 2^{-sj} < \infty \end{aligned}$$

(3) Infinite Products:

Lemma: $\{F_n\} \subseteq \Theta(n)$, $n \in \mathbb{N}$. If $\exists \{c_n\} \subseteq \mathbb{R}^+$.

St. $|F_n(z)| < c_n$, $\sum c_n < \infty$. Then

i) $\prod F_n(z) \xrightarrow{n} F(z) \in \Theta(n)$

ii) If $F_n(z) \neq 0, \forall n$. Then $\frac{F'(z)}{F(z)} = \sum \frac{F'_n(z)}{F_n(z)}$

Pf: i) $c_n \rightarrow 0 \therefore F_n(z) \neq 0$, when n is large enough.

$$\prod F_n(z) = e^{\sum \ln(1 + F_n(z))} \leq e^{\sum c_n}$$

$\therefore \prod F_n(z)$ converges.

$$\text{ii) Note that } \sum_1^{\infty} \frac{F_n'(z)}{F_n(z)} = \frac{(\pi F_n)'}{\pi F_n}$$

By i). we're done.

$$\text{e.g. } F(z) = \pi \cot \pi z = \sum_{n \in \mathbb{Z}} \frac{1}{z+n}$$

Pf: Ideal: By Liouville Thm.

prove: $A(z) = \pi \cot \pi z - \sum_{n \in \mathbb{Z}} \frac{1}{z+n}$ is bounded entire.

1) Observation:

i) $A(z+1) = A(z)$, when $z \notin \mathbb{Z}$.

ii) $F(z) = \frac{1}{z} + F_0(z)$. $F_0(z)$ is holomorphic near $z=0$.

iii) $F(z)$ has only simple isolated poles.

2) $A(z)$ is entire.

Since $z=0$ is removable by observation.

Use periodicity of $A(z)$ $\therefore z=k \in \mathbb{Z}$ are removable.

3) $A(z)$ is bounded.

using periodicity. prove $A(z)$ is bounded

in $z \in \{ |Re(z)| \leq \frac{1}{2} \}$.

Condition on $|Im(z)| \leq 1$ or $|Im(z)| > 1$.

Remark: Derive: $\frac{\sin \pi z}{\pi} = z \prod_{n \in \mathbb{Z}^+} \left(1 - \frac{z^2}{n^2} \right)$.

Note that $\left(\frac{\sin \pi z}{\pi} / z \prod_{n \in \mathbb{Z}^+} \left(1 - \frac{z^2}{n^2} \right) \right)'$

$$= \frac{\sin \pi z}{\pi} \frac{\pi \cot \pi z - \sum_{n \in \mathbb{Z}} \frac{1}{z+n}}{z \prod_{n \in \mathbb{Z}^+} \left(1 - \frac{z^2}{n^2} \right)} = 0.$$

Thm. (Weierstrass Infinite Product)

$\{a_n\} \subseteq \mathbb{C}$, $|a_n| \rightarrow \infty$ as $n \rightarrow \infty$. Then exists an entire function $f(z)$, s.t. $f(a_n) = 0, \forall n$.
 $f(z) \neq 0$, when $z \neq a_n$.

Moreover, for any other one satisfies it has form: $f(z) e^{g(z)}$, $g(z) \in \Theta(\mathbb{C})$

Pf: 1°) Canonical Function:

$$E_k(z) = (1-z) e^{\sum_{j=1}^k \frac{z^j}{a_j}}, \quad E_0(z) = 1-z$$

Lemma. For $|z| < \frac{1}{2}$, $|1 - E_k| = C|z|^{k+1}$

Pf: By $|1 - e^w| \leq |w| |e^w| \leq C|w|$.

2°) The ideal:

Insert $\prod (1 - \frac{z}{a_k})$ into $\prod E_k(z)$.

Check $f(z) = z \prod_{k=1}^{\infty} E_k(\frac{z}{a_k})$ converges.

Besides, if f_1, f_2 satisfies the condition.

Then $f_1/f_2 \in \Theta(\mathbb{C})$, nonvanishes $\therefore \frac{f_1}{f_2} = e^{g(z)}$

Remark: We have a more general Thm:

Thm. (Hadamard)

For $k \leq \ell < k+1$, $\{a_n\}$ is seq of zeros of an

entire function f . Then $f(z) = e^{p(z)} z^m \prod E_k(\frac{z}{a_k})$

m is order of zero $z=0$, $p(z)$ is a polynomial with degree $\leq k$.