

Fourier Transform in $\mathbb{C}$


Def: For $f \in \mathcal{F}_a$ if i) $f \in \mathcal{O}(\{ |Im z| < a \})$

ii) $|f(z)| \leq \frac{A}{1+x^2} \cdot \forall z = x+yi \in \{ |Im z| < a \}$.

(1) Fourier Transform on $\mathcal{F}$:

Thm. If $f \in \mathcal{F}_a$. Then $|\hat{f}(s)| \leq A e^{-2\pi b|s|} \cdot \forall 0 < b < a$.

Pf: Consider the contour:



when $s < 0$. use ①

$s \geq 0$ use ②

when $R \rightarrow \infty$. we obtain

$$\hat{f}(s) = \int_{\mathbb{R}} f(x \pm ib) e^{-2\pi i s(x \pm ib)} dx$$

Remark: For $f \in \mathcal{F}$. $\hat{f}$ has rapid decay.

Thm.

i) (Inversion Formula)

$$\text{For } f \in \mathcal{F}. f(x) = \int_{\mathbb{R}} \hat{f}(s) e^{2\pi i x s} ds.$$

ii) (Poisson Summation Formula)

$$\text{For } f \in \mathcal{F}. \sum_{n \in \mathbb{Z}} f(n) = \sum_{n \in \mathbb{Z}} \hat{f}(n)$$

Pf: i) It's from $\hat{f}(s) = \int_{\mathbb{R}} f(x \pm ib) e^{-2\pi i s(x \pm ib)} dx$.

ii) Note that $\sum \hat{f}(n) = \sum \int_{\mathbb{R}} f(x) e^{-2\pi i n x} dx$

$$= \int_{\mathbb{R}} f(x) \sum e^{-2\pi i n x} dx = \int_{\mathbb{R}} f(x) / e^{2\pi i x} dx$$

Apply Residue Formula. (only when $f \in \mathcal{F}$)

(2) Paley - Wiener Thm.

• Suppose f is valid for Inversion Transf.

Next, we will discover what condition on f will lead to $\text{supp}(\hat{f}(z)) = [-m, m]$.

Thm. If f satisfies: $|\hat{f}(z)| \leq A e^{-2\pi a|z|}$

for some $A, a > 0$. Then $f \in \mathcal{O}(S_b)$.

$$S_b = \{|z| < b\}, \quad \forall 0 < b < a.$$

Pf: $f_n(z) = \int_{-n}^n \hat{f}(z) e^{2\pi i z \eta} d\eta \in \mathcal{O}(\mathbb{C}) \rightarrow f(z)$

$$\forall z \in \bar{S}_b \subseteq S_a. \quad \therefore f(z) \in \mathcal{O}(S_b).$$

Remark: $|\hat{f}(z)| = \mathcal{O}(e^{-2\pi a|z|}) \Leftrightarrow f \in \mathcal{I}_a.$

Thm. If $f(x) \in C(\mathbb{R}) \cap M(\mathbb{R})$ Then f has an extension on $\mathbb{C}$, which is entire, satisfying: $f = \mathcal{O}(e^{2\pi m|z|})$
 $\exists m > 0, \Leftrightarrow \text{supp}(\hat{f}(z)) = [-m, m].$

Pf: ($\Leftarrow$) It's easy to estimate.

$$(\Rightarrow) 1^\circ \text{ For } f \in \mathcal{O}(\mathbb{C}), |f| \leq A \frac{e^{2\pi m|z|}}{1+x^2}$$

$$\text{By } \hat{f}(z) = \int_{\mathbb{R}} f(x+iy) e^{-2\pi i z(x+iy)} dx$$

$$\text{Then } |\hat{f}(z)| \leq C e^{-2\pi y(|z|-m)} \quad (\text{let } y \rightarrow \infty)$$

$$2^\circ \text{ For } f \in \mathcal{O}(\mathbb{C}), |f| \leq A e^{2\pi m|z|}.$$

$$\text{Approx by } f_z = f/(1+iz)^2$$

(In fact, $\frac{1}{1+x^2}$ is only for converge!)

3') Prove: $f(z)$ satisfies the condition in 2')

Lemma. $f \in \mathcal{O}(S)$, $S = \{z : -\frac{\pi}{4} < \arg z < \frac{\pi}{4}\}$, $f \in C(\bar{S})$


If $|f(z)| \leq 1$ on ∂S , $|f(z)| \leq A e^{C|z|}$ in S .

Then $|f(z)| \leq 1$, $\forall z \in \bar{S}$.

Pf: For applying MMP, we want the domain can be bounded rather than consider $z = \infty$.

Let $f_\varepsilon(z) = f(z) e^{-\varepsilon z^{\frac{1}{2}}} \in \mathcal{O}(S)$

Then $|f_\varepsilon(z)| \rightarrow 0$ (when $z \rightarrow \infty$)



We only need to consider $f_\varepsilon(z)$ in D , where $|f| \leq 1$ outside D .

Then let $\varepsilon \rightarrow 0$.

$\Rightarrow$ Next, prove: $|f(x)| \leq 1$ ($f \in M(\mathbb{R})$) $\Rightarrow |f(x+iy)| \leq A e^{B|y|}$
 $|f(z)| \leq A e^{B|z|}$

Only consider in $Q = \{x > 0, y > 0\}$. Other three quadrants remains same. Let $F = f(z) e^{22\pi i z}$

Then $|F(z)| \leq 1$ when $z \in \partial Q$, $|F| \leq A e^{B|z|}$

Rotate Q to $\{-\frac{\pi}{4} < \arg z < \frac{\pi}{4}\}$. Apply the lemma.

$\therefore |F(z)| \leq 1$ on $\bar{Q}$, i.e. $|f(z)| \leq e^{22\pi|y|}$

Thm (The case $\hat{f}(s)$ vanishes on $s < 0$)

$f(x), \hat{f}(x) \in M(\mathbb{R})$ Then $\hat{f}(s) = 0, \forall s < 0 \Leftrightarrow$

$f(x)$ can be extended continuously to $\bar{H}$, $|f| \leq M$.

$\forall z \in \bar{H}$, with $f \in \mathcal{O}(H)$

Pf: ($\Rightarrow$). By Inversion Formula, check:

$$f(z) = \int_0^\infty \bar{f(s)} e^{-2\pi i s z} ds \text{ is conti. bounded.}$$

$$\text{and approx. by } \int_0^\eta \bar{f(s)} e^{-2\pi i s z} ds \in \mathcal{O}(1)$$

$$(\Leftarrow) \text{ Consider } f_{\varepsilon, \delta}(z) = \frac{f(z+i\delta)}{(1-\varepsilon z)^2} \text{ satisfies}$$

the condition as $f(z)$. $\varepsilon, \delta > 0$.

(δ is for $f_{\varepsilon, \delta} \in \mathcal{O}(\bar{H})$. ε is for converge)

$$\text{By residue} = |\hat{f}_{\varepsilon, \delta}(s)| = \left| \int_{\mathbb{R}} f(x+i\eta) e^{2\pi i s(x+i\eta)} dx \right|$$

$$\leq \left(\int_{-\infty}^{+\infty} \frac{M}{1+x^2} dx \right) e^{2\pi s \eta} \quad \forall s < 0. \text{ Let } \eta \rightarrow \infty.$$

$$\therefore \hat{f}_{\varepsilon, \delta}(s) = 0. \quad \forall s < 0. \text{ Let } \delta \rightarrow 0. \varepsilon \rightarrow 0.$$

Prop. (Analogous conclusion for Fourier coefficients)

For $f \in \mathcal{O}(\mathcal{D}(z_0, R))$ with expansion at z_0 :

$$f(z) = \sum_{n \geq 0} a_n (z - z_0)^n. \text{ Then } a_n = \frac{1}{2\pi r^n} \int_0^{2\pi} f(z_0 + r e^{i\theta}) e^{-in\theta} d\theta$$

Pf. It's directly from $a_n = \frac{f^{(n)}(z_0)}{n!}$ by Cauchy Formula

$$\text{which also states: } 0 = \frac{1}{2\pi r^n} \int_0^{2\pi} f(z_0 + r e^{i\theta}) e^{-in\theta} d\theta$$

$\forall n < 0.$