

Vector Fields

(1) Definitions:

- A vector field on X is a Func: assigns $\forall x \in X$ to its corresponding tangent space $T_x X$.

① Vector fields:

i) For $X = U \subseteq \mathbb{R}^n$:

$$\forall x \in X. T_x U \cong \mathbb{R}^n. \text{ Def: } TU = \bigcup_{x \in U} T_x U \cong U \times \mathbb{R}^n.$$

$$\text{i.e. } TU = \{(x, v) \mid x \in U, v \in T_x U\}.$$

$$\text{Canonical Proj: } \pi: TU \rightarrow U. \quad \pi^{-1}(x) = T_x U.$$

Def: Vector field ζ on U is:

$$\begin{aligned} \zeta: U &\rightarrow TU = U \times \mathbb{R}^n. & \pi \circ \zeta &= \text{id}_U. \\ x &\mapsto (x, \tilde{\zeta}|_x) \end{aligned}$$

for some $\tilde{\zeta}|_x \in T_x U$.

ii) For arbitrary n -dim manifold X :

$$TX = \bigcup_{x \in X} T_x X = \{(x, v) \mid v \in T_x X\}. \text{ tangent bundle of } X.$$

$$\begin{aligned} \pi: TX &\rightarrow X & \pi^{-1}(x) &= T_x X. \quad \forall x \in X. \\ (x, v) &\mapsto x \end{aligned}$$

Remark: TX may not be cross-product of X with some other set.

It naturally has structure of manifold.

$$\text{so, } \dim(TX) = 2 \dim(X).$$

Def: Vector field on X is Func:

$$\zeta: X \rightarrow TX, \quad \pi \circ \zeta = 1_X$$

② Smooth Structure:

i) For $\tilde{U} \subseteq_{\text{open}} \mathbb{R}^n$, $\tilde{\zeta}: \tilde{U} \rightarrow T\tilde{U}$.

It's surely that $\tilde{\zeta}$ is smooth $\Leftrightarrow$ the second component of $\tilde{\zeta}$ is smooth.

ii) For n -dim manifold X , $(U, f) \in \mathcal{A}_X$, ζ is vector field

$$TU = \pi^{-1}(U) \subseteq TX. \text{ See it in chart by:}$$

$$F: TU \xrightarrow{\cong} T\tilde{U} = \tilde{U} \times \mathbb{R}^n$$

$$(x, v) \mapsto (f(x), Af(v))$$

$$\Rightarrow F \circ \zeta|_U \circ f^{-1}: \tilde{U} \rightarrow T\tilde{U} \cong \tilde{U} \times \mathbb{R}^n$$

$$f(x) \mapsto (f(x), Af \circ \zeta|_x \circ f^{-1}(f(x)))$$

$$\text{i.e. } F \circ \zeta|_U \circ f^{-1} = (1_{\tilde{U}}, Af \circ \zeta|_x \circ f^{-1}) \triangleq (1_{\tilde{U}}, \tilde{\zeta})$$

$$\tilde{\zeta}: \tilde{U} \rightarrow \mathbb{R}^n$$

Remark: So TX locally looks like an open subset of $\mathbb{R}^{2n}$.

Def: A vector field $\zeta: X \rightarrow TX$ is smooth

$$\text{if } \forall (U, f) \in \mathcal{A}_X, F \circ \zeta|_U \circ f^{-1} = (1_{\tilde{U}}, \tilde{\zeta}):$$

$$\tilde{U} \rightarrow \tilde{U} \times \mathbb{R}^n, \text{ its } 2^{\text{nd}}\text{-component is smooth.}$$

Remark: It's inept with the choice of charts:

$$\text{For: } \tilde{F}_1: F_1 \circ g|_U \circ f_1^{-1} = (1_{\tilde{U}}, \tilde{f}_1): \tilde{U} \rightarrow \tilde{U} \times \mathbb{R}^n$$

$$\tilde{F}_2: F_2 \circ g|_U \circ f_2^{-1} = (1_{\tilde{U}}, \tilde{f}_2): \tilde{U} \rightarrow \tilde{U} \times \mathbb{R}^n.$$

$U = U_1 \cap U_2$, Next, we define:

$$\phi_{21}: \tilde{U} \times \mathbb{R}^n \xrightarrow{\sim} \tilde{U} \times \mathbb{R}^n$$

$$(\tilde{x}, \tilde{v}) \mapsto (\phi_{21}(\tilde{x}), D\phi_{21}|_{\tilde{x}} \cdot \tilde{v})$$

$$\therefore \tilde{F}_2 = \phi_{21} \circ \tilde{F}_1. \text{ Besides: } \tilde{f}_2|_{\phi_{21}(\tilde{x})} = D\phi_{21}|_{\tilde{x}} \cdot \tilde{f}_1|_{\tilde{x}}$$

(2) Vector Fields from

Translation Law:

Def: $\mathcal{F}$ is a rule, st. $\mathcal{F}: (U, f) \mapsto \tilde{f}_f, \forall (U, f) \in A_X$.

$\tilde{f}_f: \tilde{U} \rightarrow \mathbb{R}^n$ smooth. Besides, $\forall (U_1, f_1), (U_2, f_2) \in A_X$,

$$\forall \tilde{x} \in f_1(U_1 \cap U_2), \tilde{f}_2|_{\phi_{21}(\tilde{x})} = D\phi_{21}|_{\tilde{x}} \cdot \tilde{f}_1|_{\tilde{x}}$$

prop. $\mathcal{F}$ defines a smooth vector field on X .

Pf: Restrict $\mathcal{F}$ on A_X^x :

$$\mathcal{F}_x = \mathcal{F}|_{A_X^x}: A_X^x \longrightarrow \mathbb{R}^n \quad \text{this is a tangent vector}$$

$$(U_x, f_x) \longmapsto \tilde{f}_f|_{f_x}$$

$$\therefore \tilde{\mathcal{F}}: X \longrightarrow TX \quad \text{is a smooth vector field.}$$

$$x \longmapsto (x, \mathcal{F}_x)$$

since $\tilde{f}_f|_{f_x}$ is smooth, $\mathcal{F}$ corresponds to $\mathcal{F}_x, (U_x, f_x)$

(3) Flow:

① Def. A Flow on X is a smooth map:

$$F: (-\varepsilon, \varepsilon) \times X \rightarrow X. \text{ Besides, } \forall s \in (-\varepsilon, \varepsilon), \text{ fixed.}$$

$$F_s(x) = X \rightarrow X \text{ is } C^\infty \text{-diffeomorphism. } F_0 = \text{Id}_X.$$

Remark: $(-\varepsilon, \varepsilon) \times X$ is naturally a smooth manifold

with dimension: $\dim X + 1$. $\hookrightarrow$ Chart:

$$(1 \times F, (-\varepsilon, \varepsilon) \times U).$$

e.g. $F_s(x) = T' \rightarrow T'$ is flow of T' .
 $[x] \mapsto [x+s]$

$\Rightarrow$ If we fix $x \in X$, $F_x(s) = (-\varepsilon, \varepsilon) \rightarrow X$.

$F_x(0) = F(0, x) = x$, it's a curve through x .

Repeat on all $x \in X$. We obtain a vector field

$$\text{on } X: \mathcal{F}^F: X \rightarrow (TX) \times \{F_x(s)\}_{s \in (-\varepsilon, \varepsilon)}.$$

Remark: We can view vector field as the infinitesimal version of flow.

② Procedure in charts:

$$\tilde{F} = f \circ F \circ (1 \times f)^{-1} = (\tilde{F}_1, \tilde{F}_2, \dots, \tilde{F}_n) : (-\varepsilon, \varepsilon) \times \tilde{U} \rightarrow \tilde{U}$$

and $\tilde{F}(0, x) = \text{Id}_{\tilde{U}}$. The associated vector field is:

$$\tilde{\mathcal{F}}^F = \frac{\partial \tilde{F}}{\partial s} \Big|_{s=0} = \left(\frac{\partial \tilde{F}_1}{\partial s} \Big|_{s=0}, \dots, \frac{\partial \tilde{F}_n}{\partial s} \Big|_{s=0} \right) : \tilde{U} \rightarrow \mathbb{R}^n.$$

③ Existence:

If we have vector field $\mathcal{F}$ on X , cpt manifold. Then we

can find flow F , st. $\mathcal{F}^F = \mathcal{F}$. If X is compact, ε may be $\rightarrow \infty$.